


$D_0(2550)^0$ $I(J^P) = \frac{1}{2}(0^-)$

OMMITTED FROM SUMMARY TABLE
 $J^P = 0^-$ determined by AAIJ 20D.

 $D_0(2550)^0$ MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2549 ± 19 OUR AVERAGE				Error includes scale factor of 4.6. See the ideogram below.
2518 ± 2 ± 7	79k	¹ AAIJ	20D LHCb	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2579.5 ± 3.4 ± 5.5	60k	AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$
2539.4 ± 4.5 ± 6.8	34k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$



¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.
 $D_0(2550)^0$ MASS (MeV)

 $D_0(2550)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
165 ± 24 OUR AVERAGE				Error includes scale factor of 2.0. See the ideogram below.
199 ± 5 ± 17	79k	¹ AAIJ	20D LHCb	$B^- \rightarrow D^{*+} \pi^- \pi^-$
177.5 ± 17.8 ± 46.0	60k	AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$
130 ± 12 ± 13	34k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$



¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.
 $D_0(2550)^0$ WIDTH (MeV)

$D_0(2550)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad D^{*+} \pi^-$	seen

$D_0(2550)^0$ POLARIZATION AMPLITUDE A_{D_J}

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$ decays.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 1.3	60k	¹ AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$

¹ Systematic uncertainty not estimated.

$D_0(2550)^0$ REFERENCES

AAIJ 20D PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ 13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA... 10P PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)